

REFERENCES

- [1] “Industrial Pollution Control Guidelines: No. 3 - Desiccated Coconut Industry,” 1992.
- [2] D. G. Cirne, X. Paloumet, L. Björnsson, M. M. Alves, and B. Mattiasson, “Anaerobic digestion of lipid-rich waste-Effects of lipid concentration,” *Renew. Energy*, vol. 32, no. 6, pp. 965–975, 2007.
- [3] C. Ye, C. Jay J, and C. Kurt S., “Inhibition of anaerobic digestion process: A review Ye,” *Bioresour. Technol.*, vol. 99, pp. 4044–4064, 2007.
- [4] N. Rasit, A. Idris, R. Harun, W. Azlina, W. Ab, and K. Ghani, “Effects of lipid inhibition on biogas production of anaerobic digestion from oily effluents and sludges: An overview,” *Renew. Sustain. Energy Rev.*, vol. 45, pp. 351–358, 2015.
- [5] Carlos Augusto de Lemos Chernicharo, *Anaerobic Reactors*. IWA Publishing, 2007.
- [6] B. K. T. Samarasiri, P. A. D. Mihiranga, and P. G. Rathnasiri, “Effect of Lipid Inhibition in Anaerobic Wastewater Treatment: A Case Study using Desiccated Coconut Wastewater Effect of Lipid Inhibition in Anaerobic Wastewater Treatment: A Case Study using Desiccated Coconut,” in *Annual Sessions of IESL*, 2016, no. October, pp. 295–304.
- [7] M. & E. Inc., G. Tchobanoglous, F. L. Burton, and H. D. Stensel, *Wastewater Engineering: Treatment and Reuse*, 4th Editio. McGraw Hill Higher Education, 2003.
- [8] H. N. Chanakya, H. K. Khuntia, N. Mukherjee, R. Aniruddha, J. R. Mudakavi, and P. Thimmaraju, “The physicochemical characteristics and anaerobic degradability of desiccated coconut industry waste water,” *Environ. Monit. Assess.*, vol. 187, no. 12, p. 772, Dec. 2015.
- [9] D. R. Rosa *et al.*, “Performance and molecular evaluation of an anaerobic system with suspended biomass for treating wastewater with high fat content after enzymatic hydrolysis,” *Bioresour. Technol.*, 2009.
- [10] A. A. Mendes, E. B. Pereira, and H. F. de Castro, “Effect of the enzymatic hydrolysis pretreatment of lipids-rich wastewater on the anaerobic biodigestion,” *Biochem. Eng. J.*, vol. 32, no. 3, pp. 185–190, Dec. 2006.
- [11] S. Cherif, F. Aloui, F. Carrière, and S. Sayadi, “Lipase Pre-Hydrolysis Enhance Anaerobic Biodigestion of Soap Stock from an Oil Refining Industry,” vol. 114, no. 2, pp. 109–114, 2014.
- [12] P. Taylor, F. R. C. Damasceno, D. M. G. Freire, and M. C. Cammarota,

“Assessing a mixture of biosurfactant and enzyme pools in the anaerobic biological treatment of wastewater with a high-fat content,” no. December 2014, pp. 37–41.

- [13] K. Sri, B. Kameswari, C. Kalyanaraman, S. Porselvam, K. Thanasekaran, and E. T. Division, “Enhancement of Biogas Generation by Addition of Lipase in the Co-Digestion of Tannery Solid Wastes,” vol. 39, no. 8, pp. 781–786, 2011.
- [14] I. Angelidaki, S. P. Petersen, and B. K. Ahring, “Effects of lipids on thermophilic anaerobic digestion and reduction of lipid inhibition upon addition of bentonite,” *Appl. Microbiol. Biotechnol.*, 1990.
- [15] E. Nordell, A. B. Hansson, and M. Karlsson, “Zeolites relieves inhibitory stress from high concentrations of long chain fatty acids,” *Waste Manag.*, vol. 33, no. 12, pp. 2659–2663, Dec. 2013.
- [16] T. V. Adulkar and V. K. Rathod, “Ultrasound assisted enzymatic pre-treatment of high fat content dairy wastewater,” *Ultrason. Sonochem.*, 2014.
- [17] N. A. Oz and C. C. Yarimtepe, “Ultrasound assisted biogas production from landfill leachate,” *Waste Manag.*, vol. 34, no. 7, pp. 1165–1170, Jul. 2014.
- [18] L.-J. Wu, T. Kobayashi, Y.-Y. Li, and K.-Q. Xu, “Comparison of single-stage and temperature-phased two-stage anaerobic digestion of oily food waste,” *Energy Convers. Manag.*, vol. 106, pp. 1174–1182, Dec. 2015.
- [19] M. BECCARI, M. MAJONE, and L. TORRISI, “Two-reactor system with partial phase separation for anaerobic treatment of olive oil mill effluents,” *Water Sci. Technol.*, vol. 38, no. 4–5, pp. 53–60, 1998.
- [20] S. Astals, D. J. Batstone, J. Mata-Alvarez, and P. D. Jensen, “Identification of synergistic impacts during anaerobic co-digestion of organic wastes,” *Bioresour. Technol.*, 2014.
- [21] L. Martín-González, L. F. Colturato, X. Font, and T. Vicent, “Anaerobic co-digestion of the organic fraction of municipal solid waste with FOG waste from a sewage treatment plant: Recovering a wasted methane potential and enhancing the biogas yield,” *Waste Manag.*, vol. 30, no. 10, pp. 1854–1859, Oct. 2010.
- [22] R. Affes, J. Palatsi, X. Flotats, H. Carrère, J. P. Steyer, and A. Battimelli, “Saponification pretreatment and solids recirculation as a new anaerobic process for the treatment of slaughterhouse waste,” *Bioresour. Technol.*, 2013.
- [23] R. Uma Rani, S. Kaliappan, S. Adish Kumar, and J. Rajesh Banu, “Combined treatment of alkaline and disperser for improving solubilization and anaerobic biodegradability of dairy waste activated sludge,” *Bioresour. Technol.*, vol. 126, pp. 107–116, Dec. 2012.

- [24] M. López Torres and M. d C. Espinosa Llorens, “Effect of alkaline pretreatment on anaerobic digestion of solid wastes,” *Waste Manag.*, 2008.
- [25] J. M. Berg, J. L. Tymoczko, and L. Stryer, *Biochemistry*, 7th Editio. Palgrave MacMillan, 2011.
- [26] R. F. Domingues *et al.*, “Effect of enzymatic pretreatment on the anaerobic digestion of milk fat for biogas production,” *Food Res. Int.*, vol. 73, pp. 26–30, Jul. 2015.
- [27] Y. Meng, F. Luan, H. Yuan, X. Chen, and X. Li, “Enhancing anaerobic digestion performance of crude lipid in food waste by enzymatic pretreatment,” *Bioresour. Technol.*, 2016.
- [28] P. Taylor *et al.*, “The effect of enzymatic pre-hydrolysis of dairy wastewater on the granular and immobilized microbial community in anaerobic bioreactors,” no. December, pp. 37–41, 2013.
- [29] E. Efremenko, O. Senko, D. Zubaerova, E. Podorozhko, and V. Lozinsky, “New Biocatalyst with Multiple Enzymatic Activities for Treatment of Complex Food Wastewaters,” vol. 9862, no. 2, pp. 208–212, 2008.
- [30] V. M. F. Alexandre, A. M. Valente, M. C. Cammarota, and D. M. G. Freire, “Performance of anaerobic bioreactor treating fish-processing plant wastewater pre-hydrolyzed with a solid enzyme pool,” *Renew. Energy*, vol. 36, no. 12, pp. 3439–3444, 2011.
- [31] K. J. Kennedy and J. D. B. O, “Effect of enzymes on anaerobic digestion of primary sludge and septic tank performance,” 2012.
- [32] A. B. G. Valladão, D. M. G. Freire, and M. C. Cammarota, “Enzymatic pre-hydrolysis applied to the anaerobic treatment of effluents from poultry slaughterhouses,” *Int. Biodeterior. Biodegrad.*, 2007.
- [33] G. Dors and A. A. Mendes, “Simultaneous enzymatic hydrolysis and anaerobic biodegradation of lipid-rich wastewater from poultry industry,” pp. 343–349, 2013.
- [34] E. Mobarak-Qamsari, R. Kasra-Kermanshahi, M. Nosrati, and T. Amani, “Enzymatic pre-hydrolysis of high fat content dairy wastewater as a pretreatment for anaerobic digestion,” *Int. J. Environ. Res.*, vol. 6, no. 2, pp. 475–480, 2012.
- [35] Z. Ning, J. Ji, Y. He, Y. Huang, G. Liu, and C. Chen, “E ff ect of Lipase Hydrolysis on Biomethane Production from Swine Slaughterhouse Waste in China,” 2016.
- [36] *Oil and Grease, Hexane Extractable Gravimetric Method*. U. S. Environmental Protection Agency.

- [37] R. Uma Rani, S. Adish Kumar, S. Kaliappan, I. T. Yeom, and J. Rajesh Banu, "Low temperature thermo-chemical pretreatment of dairy waste activated sludge for anaerobic digestion process," *Bioresour. Technol.*, vol. 103, no. 1, pp. 415–424, 2012.
- [38] K. S. B. Kameswari, C. Kalyanaraman, S. Porselvam, and K. Thanasekaran, "Optimization of inoculum to substrate ratio for bio-energy generation in co-digestion of tannery solid wastes," pp. 241–250, 2012.
- [39] A. A. Mendes, E. B. Pereira, and A. F. Jr, "Anaerobic Biodegradability of Dairy Wastewater Pretreated with Porcine Pancreas Lipase," vol. 53, no. December, pp. 1279–1284, 2010.