

## REFERENCES

- Abderrahim, B., Abderrahman, E., Mohamed, A., Fatima, T., Abdesselam, T., & Krim, O. (2015). Kinetic Thermal Degradation of Cellulose, Polybutylene Succinate and a Green Composite: Comparative Study. *World Journal of Environmental Engineering*, 95-110. doi:10.12691/wjee-3-4-1
- Abuzerr, S., Darwish, M., & Mahvi, A. H. (2018, April). Simultaneous removal of cationic methylene blue and anionic reactive red 198 dyes using magnetic activated carbon nanoparticles: equilibrium, and kinetics analysis. (W. Rauch, Ed.) *Water Science Techonology*, 2017(2), 534–545. doi:10.2166/wst.2018.145
- Aguilar, Z. P. (2013). Nanomaterials for Medical Applications. In *Types of Nanomaterials and Corresponding Methods of Synthesis* (pp. 33-82). doi:10.1016/B978-0-12-385089-8.00002-9
- Ahmad, A., Rafatullah, M., Sulaiman, O., Ibrahim, M. H., & Hashim, R. (2009, October). Scavenging behaviour of meranti sawdust in the removal of methylene blue from aqueous solution. (Z. (. He, Ed.) *Journal of Hazardous Materials*, 170(1), 357-365. doi:10.1016/j.jhazmat.2009.04.087
- Al-Degs, Y., El-Barghouthi, M., El-Sheikh, A., & Walker, G. (2008). Effect of solution pH, ionic strength, and temperature on adsorption behavior of reactive dyes on activated carbon. *Dyes and Pigments*, 16-23. doi:10.1016/j.dyepig.2007.03.001
- Almeida, S. L., Schmidt, É. C., Pereira, D. T., Kreusch, M., Felix, M. R., Osorio, L. K., Bouzon, Z. L. (2013, May). Effect of ultraviolet-B radiation in laboratory on morphological and ultrastructural characteristics and physiological parameters of selected cultivar of *Oryza sativa* L. doi:10.1007/s00709-013-0512-0
- Alshamsi, F. A., Albadwawi, A. S., Alnuaimi, M. M., Rauf, M. A., & Ashraf, S. S. (2007). Comparative efficiencies of the degradation of Crystal Violet using

UV/hydrogen peroxide and Fenton's reagent. (B. M. Heron, & M. Wainwright, Eds.) *Dyes and Pigments*, 74(2), 283-287. doi:10.1016/j.dyepig.2006.02.016

Amarasinghe, B., & Amarasinghe, A. (2020, October). Study on Mass Transfer, Kinetic Parameters and Rate Determining Step with Statistical Analysis in Adsorption of Pb ions onto Coir Pith. *International Journal of Scientific Research and Innovative Technology*, 7(9). Retrieved from [https://ijsrit.com/uploaded\\_all\\_files/2494925667\\_f4.pdf](https://ijsrit.com/uploaded_all_files/2494925667_f4.pdf)

Amarasinghe, P., & Amarasinghe, K. (2020). Determination of Mass Transfer Coefficients for Adsorption of Pb and Cd Onto Coir Pith and Statistical Analysis. *Fortune journals*, 2(3), 139. doi:DOI: 10.26502/jatri.017

Amin, M. T., Alazba, A. A., & Shafiq, M. (2015, November). Adsorptive Removal of Reactive Black 5 from Wastewater Using Bentonite Clay: Isotherms, Kinetics and Thermodynamics. (M. A. Rosen, Ed.) *Sustainability*, 7(11), 15302-15318. doi:10.3390/su71115302

Ayed, L., Cheriaa, J., Laadhari, N., Cheref, A., & Bakhrouf, A. (2009). Biodegradation of crystal violet by an isolated *Bacillus* sp. (F. Cappitelli, Ed.) *Annals of Microbiology*, 59(2), 267-272. doi:10.1007/BF03178327

Baghapour, A. M., Mahvi, A. H., & Pourfadakari, S. (2013). Thermodynamic analysis of reactive red 198 removal from synthetic wastewater by using multiwall carbon nanotubes. (R. Mirzaei, Ed.) *Health Scope*, 2(3), 149-155. Retrieved from <https://www.sid.ir/en/Journal/ViewPaper.aspx?ID=437199>

Bahmani, P., Kalantary, R. R., Esrafil, A., Gholami, M., & Jafari, A. J. (2013, June). Evaluation of Fenton oxidation process coupled with biological treatment for the removal of reactive black 5 from aqueous solution. (S. Nasser, Ed.) *Journal of Environmental Health Science and Engineering*, 11(1). doi:10.1186/2052-336X-11-13

Banat, F., Al-Asheh, S., Al-Anbar, S., & Al-Refai, S. (2006, June). Microwave- and acid-treated bentonite as adsorbents of methylene blue from a simulated

dye wastewater. (L. Wong, & A. Basu, Eds.) *Bulletin of Engineering Geology and the Environment*, 66(1), 53-58. doi:10.1007/s10064-006-0054-1

Baneshi, M., Naraghi, B., Rahdar, S., Biglari, H., Saeidi, M., Ahamadabadi, M., Alipour, V. (2016). Removal of remazol black B dye from aqueous solution by electrocoagulation equipped with iron and aluminium electrodes. (V. Azevedo, Ed.) *The IIOAB Journal*, 7, 529-535. Retrieved from [https://www.iioab.org/articles/IIOABJ\\_7.S2\\_529-535.pdf](https://www.iioab.org/articles/IIOABJ_7.S2_529-535.pdf)

Bazzo, A., Adebayo, M. A., Dias, S. L., Lima, E. C., Vagheti, J. C., & de Oliveira, E. R. (2015, January). Avocado seed powder: characterization and its application for crystal violet dye removal from aqueous solutions. *Desalination and Water Treatment*, 57(34), 15873-15888. doi:10.1080/19443994.2015.1074621

Belhachemi, M., Belala, Z., Lahcene, D., & Addoun, F. (2009). Adsorption of phenol and dye from aqueous solution using chemically modified date pits activated carbons. *Desalination and Water Treatment*, 182-190. doi:10.5004/dwt.2009.729

Berradi, M., Hsissou, R., Khudhair, M., Assouag, M., Cherkaoui, O., Bachiri, A. E., & Harfi, A. E. (2019, November 05). Textile finishing dyes and their impact on aquatic environs. *Heliyon*, 05(11). doi:10.1016/j.heliyon.2019.e02711

Bhattacharyya, K. G., & Sharma, A. (2005, April). Kinetics and thermodynamics of Methylene Blue adsorption on Neem (*Azadirachta indica*) leaf powder. (B. M. Heron, & M. Wainwright, Eds.) *Dyes and Pigments*, 65(1), 51-59. doi:10.1016/j.dyepig.2004.06.016

Bhaumik, M., McCrindle, R. I., Maity, A., Agarwal, S., & Gupta, V. K. (2016, March). Polyaniline nanofibers as highly effective re-usable adsorbent for removal of reactive black 5 from aqueous solutions. (M. Malmsten, Ed.) *Journal of Colloid and Interface Science*, 466, 442-451. doi:10.1016/j.jcis.2015.12.056

- Bouraie, M. E., & Din, W. S. (2016, September). Biodegradation of Reactive Black 5 by *Aeromonas hydrophila* strain isolated from dye-contaminated textile wastewater. *Sustainable Environment Research*, 26(5), 209-216. doi:10.1016/j.serj.2016.04.014
- Bouras, H. D., Yeddou, A. R., Bouras, N., Hellel, D., Holtz, M. D., Sabaou, N., Nadjemi, B. (2017, November). Biosorption of Congo red dye by *Aspergillus carbonarius* M333 and *Penicillium glabrum* Pg1: Kinetics, equilibrium and thermodynamic studies. (H.-K. Tsao, Ed.) *Journal of the Taiwan Institute of Chemical Engineers*, 80, 915-923. doi:10.1016/j.jtice.2017.08.002
- Brito, S. M., Andrade, H. M., Soares, L. F., & Azevedo, R. P. (2010, February). Brazil nut shells as a new biosorbent to remove methylene blue and indigo carmine from aqueous solutions. (D. Aga, A. J. Daugulis, & Z. (. He, Eds.) *Journal of Hazardous Materials*, 174(1-3), 84-92. doi:10.1016/j.jhazmat.2009.09.020
- Bunaciu, A. A., Fleschin, Ş., & Aboul-Enein, H. Y. (2013). Evaluation of the Protein Secondary Structures Using Fourier Transform Infrared Spectroscopy. *Gazi University Journal of Science*, 27(1), 637-644. Retrieved from [https://www.researchgate.net/publication/286655543\\_Evaluation\\_of\\_the\\_protein\\_secondary\\_structures\\_using\\_fourier\\_transform\\_infrared\\_spectroscopy](https://www.researchgate.net/publication/286655543_Evaluation_of_the_protein_secondary_structures_using_fourier_transform_infrared_spectroscopy)
- Cardoso, N. F., Lima, E. C., Calvete, T., Pinto, I. S., Amavisca, C. V., Fernandes, T. H., Alencar, W. S. (2011, March). Application of Aqai Stalks As Biosorbents for the Removal of the Dyes Reactive Black 5 and Reactive Orange 16 from Aqueous Solution. (J. Siepmann, Ed.) *Journal of Chemical & Engineering Data*, 56(5), 1857–1868. doi:10.1021/je100866c
- Carneiro, P. A., Umbuzeiro, G. A., Oliveira, D. P., & Zanoni, M. V. (2010, February). Assessment of water contamination caused by a mutagenic textile. (D. Aga, A. J. Daugulis, G. Lyberatos, & Z. He, Eds.) *Journal of Hazardous Materials*, 174(1-3), 694-699. doi:10.1016/j.jhazmat.2009.09.106

- Çelebi, H. (2019, January). The applicability of evaluable wastes for the adsorption of Reactive Black 5. (M. Abbaspour, Ed.) *International Journal of Environmental Science and Technology*, 16(1), 135-146. doi:10.1007/s13762-018-1969-3
- Chakraborty, S., Basu, J. K., De, S., & DasGupta, S. (2006, May). Adsorption of Reactive Dyes from a Textile Effluent Using Sawdust as the Adsorbent. (P. E. Savage, Ed.) *Industrial & Engineering Chemistry Research*, 45(13), 4732–4741. doi:10.1021/ie050302f
- Chakraborty, S., Mukherjee, A., Das, S., Maddela, N. R., Iram, S., & Das, P. (2021, February). Study on isotherm, kinetics, and thermodynamics of adsorption of crystal violet dye by calcium oxide modified fly ash. (C. Kwang-Ho, Ed.) *Environmental Engineering Research*, 26(1). doi:10.4491/eer.2019.372
- Chan, S.-L., Tan, Y., Abdullah, A., & Ong, S.-T. (2016, April). Equilibrium, kinetic and thermodynamic studies of a new potential biosorbent for the removal of Basic Blue 3 and Congo Red dyes: Pineapple (*Ananas comosus*) plant stem. (H.-K. Tsao, Ed.) *Journal of the Taiwan Institute of Chemical Engineers*, 61, 306-3015. doi:10.1016/j.jtice.2016.01.010
- Chaukura, N., Murimba, E. C., & Gwenzzi, W. (2017, November). Sorptive removal of methylene blue from simulated wastewater using biochars derived from pulp and paper sludge. (R. Naidu, D. L. Nghiem, & M. Rood, Eds.) *Environmental Technology & Innovation*, 8, 132-140. doi:10.1016/j.eti.2017.06.004
- Chawla, S., Uppal, H., Yadav, M., Bahadur, N., & Singh, N. (2017, January). Zinc peroxide nanomaterial as an adsorbent for removal of Congo red dye from waste water. (R. Handy, & B. Yan, Eds.) *Ecotoxicology and Environmental Safety*, 135, 68-74. doi:10.1016/j.ecoenv.2016.09.017
- Cheng, S., Zhang, L., Ma, A., Xia, H., Peng, J., Li, C., & Shu, J. (2018, March). Comparison of activated carbon and iron/cerium modified activated carbon to

- remove methylene blue from wastewater. (J. Hao, & X. C. Le, Eds.) *Journal of Environmental Sciences*, 65, 92-102. doi:10.1016/j.jes.2016.12.027
- Cheruiyot, G. K., Wanyonyi, W. C., Kiplimo, J. J., & Maina, E. N. (2019, September). Adsorption of toxic crystal violet dye using coffee husks: Equilibrium, kinetics and thermodynamics study. (B. Gyampoh, Ed.) *Scientific African*, 5. doi:10.1016/j.sciaf.2019.e00116
- Chong, M., Cho, Y., Poh, P., & Jin, B. (2015, February). Evaluation of Titanium dioxide photocatalytic technology for the treatment of reactive Black 5 dye in synthetic and real greywater effluents. (J. J. Klemeš, C. M. de Almeida, & Y. Wang, Eds.) *Journal of Cleaner Production*, 89, 196-202. doi:10.1016/j.jclepro.2014.11.014
- Croce, R., Cina, F., Lombardo, A., Crispeyn, G., Cappelli, C. I., Vian, M., Baderna, D. (2017, October). Aquatic toxicity of several textile dye formulations: Acute and chronic assays with *Daphnia magna* and *Raphidocelis subcapitata*. (R. Handy, & B. Yan, Eds.) *Ecotoxicology and Environmental Safety*, 144, 79-87. doi:10.1016/j.ecoenv.2017.05.046
- Dahr, M. F., Abolghasemi, H., Esmaili, M., Shojamoradi, A., & Fatoorehchi, H. (2014, September). Adsorption Characteristics of Congo Red from Aqueous Solution onto Tea Waste. (A. Gill, Ed.) *Chemical Engineering Communications*, 202(2), 181-193. doi:10.1080/00986445.2013.836633
- Dbik, A., Bentahar, S., Khomri, M. E., Messaoudi, N. E., & Lacherai, A. (2020). Adsorption of Congo red dye from aqueous solutions using tunics of the corm of the saffron. *Materials Today : Proceedings*, 22, 134-139. doi:10.1016/j.matpr.2019.08.148
- Dehghani, M., Shahsavani, S., Jamshidi, F., & Shamsedini, N. (2019). Comparison of the Efficiency of Fenton and Photo-Fenton Processes for the Removal of Reactive Red 198 Dye from the Aqueous Solution. *Iranian Journal of Health, Safety & Environment*, 6(4), 1336-1342. Retrieved from <https://www.ijhse.ir/index.php/IJHSE/article/view/419>

- Deive, F. J., Domínguez, A., Barrio, T., Moscoso, F., Morán, P., Longo, M. A., & Sanromán, M. A. (2010, October). Decolorization of dye Reactive Black 5 by newly isolated thermophilic microorganisms from geothermal sites in Galicia (Spain). (D. Aga, A. J. Daugulis, & Z. He, Eds.) *Journal of Hazardous Materials*, 182(1-3), 735-742. doi:10.1016/j.jhazmat.2010.06.096
- Elkady, M. F., Ibrahim, A. M., & AbdEl-Latif, M. M. (2011, September). Assessment of the adsorption kinetics, equilibrium and thermodynamic for the potential removal of reactive red dye using eggshell biocomposite beads. (N. Hilal, Ed.) *Desalination*, 278(1-3), 412-423. doi:10.1016/j.desal.2011.05.063
- Elsherif, K. M., El-Dali, A., Alkarewi, A. A., Ewlad-Ahmed, A. M., & Treban, A. (2021). Adsorption of Crystal Violet Dye Onto Olive Leaves Powder: Equilibrium and Kinetic Studies. *Chemistry International*, 7(2), 79-89. doi:10.5281/zenodo.4441851
- Eren, Z., & Acar, F. N. (2006, June). Adsorption of Reactive Black 5 from an aqueous solution: equilibrium and kinetic studies. (N. Hilal, Ed.) *Desalination*, 194(1-3), 1-10. doi:10.1016/j.desal.2005.10.022
- Eslami, H., Shariatifar, A., Rafiee, E., Shiranian, M., Salehi, F., Hosseini, S. S., Ebrahimi, A. A. (2019, February). Decolorization and biodegradation of reactive Red 198 Azo dye by a new *Enterococcus faecalis*–*Klebsiella variicola* bacterial consortium isolated from textile wastewater sludge. (P. J. Large, Ed.) *World Journal of Microbiology and Biotechnology*, 35. doi:10.1007/s11274-019-2608-y
- Esmaeili, A., & Kalantari, M. (2015, February). Bioremoval of an azo textile dye, Reactive Red 198, by *Rhizopus oryzae*. *Desalination and Water Treatment*, 57(14), 6401-6410. doi:10.1080/19443994.2015.1010237
- Fabryanty, R., Valencia, C., Soetaredjo, F. E., Putro, J. N., Santoso, S. P., Kurniawan, A., Ismadji, S. (2017, December). Removal of crystal violet dye by adsorption using bentonite – alginate composite. (G. L. Dotto, D. Fatta-

- Kassinis, & Y. Lee, Eds.) *Journal of Environmental Chemical Engineering*, 5(6), 5677-5687. doi:10.1016/j.jece.2017.10.057
- Gedam, V. V., Raut, P., Chahande, A., & Pathak, P. (2019, March). Kinetic, thermodynamics and equilibrium studies on the removal of Congo red dye using activated teak leaf powder. (E. Drioli, & S. A. Aljlil, Eds.) *Applied Water Science*, 9(3). doi:10.1007/s13201-019-0933-9
- Ghaneian, M. T., Jamshidi, B., Dehvari, M., & Amrollahi, M. (2015, May). Pomegranate seed powder as a new biosorbent of reactive red 198 dye from aqueous solutions: adsorption equilibrium and kinetic studies. (M. Anpo, Ed.) *Research on Chemical Intermediates*, 41(5), 3223–3234. doi:10.1007/s11164-013-1427-2
- Gita, S., Hussan, A., & Choudhury, T. G. (2017, December). Impact of Textile Dyes Waste on Aquatic Environments and its Treatment. (S. K. Konar, Ed.) *Environment & Ecology*, 35(3C), 2349-2353. Retrieved from [https://www.researchgate.net/publication/321443064\\_Impact\\_of\\_Textile\\_Dyes\\_Waste\\_on\\_Aquatic\\_Environments\\_and\\_its\\_Treatment](https://www.researchgate.net/publication/321443064_Impact_of_Textile_Dyes_Waste_on_Aquatic_Environments_and_its_Treatment)
- Grassi, P., Drumm, F. C., Georgin, J., Franco, D. S., Dotto, G. L., Foletto, E. L., & Jahn, S. L. (2020, October). Application of Cordia trichotoma sawdust as an effective biosorbent for removal of crystal violet from aqueous solution in batch system and fixed-bed column. (P. Garrigues, Ed.) *Environmental Science and Pollution Research*, 6771–6783. doi:<https://doi.org/10.1007/s11356-020-11005-6>
- Grassi, P., Reis, C., Drumm, F. C., Georgin, J., Tonato, D., Escudero, L. B., Dotto, G. L. (2019, February). Biosorption of crystal violet dye using inactive biomass of the fungus Diaporthe schini. (W. Rauch, Ed.) *Water Science & Technology*, 79(4), 709-717. doi:10.2166/wst.2019.091
- Gül, Ü. D. (2013, October). Treatment of dyeing wastewater including reactive dyes. *WaterSA*, 39(5). doi:10.4314/wsa.v39i5.2



- Gulnaz, O., Sahmurova, A., & Kama, S. (2011, Nov). Removal of Reactive Red 198 from aqueous solution by *Potamogeton crispus*. (S. Allen, T. Aminabhavi, & D. Dionysiou, Eds.) *Chemical Engineering Journal*, 174(2-3), 579-585. doi:10.1016/j.cej.2011.09.061
- Gupta, V. K., Pathania, D., Kothiyal, N. C., & Sharma, G. (2014, February). Polyaniline zirconium (IV) silicophosphate nanocomposite for remediation of methylene blue dye from waste water. (W. Schröer, T. Yamaguchi, & A. J. Valente, Eds.) *Journal of Molecular Liquids*, 190, 139-145. doi:10.1016/j.molliq.2013.10.027
- Haffad, H., Zbair, M., Anfar, Z., Ahsaine, H. A., Bouhlal, H., & Khallok, H. (2019, March). Removal of reactive red-198 dye using chitosan as an adsorbent: optimization by Central composite design coupled with response surface methodology. (R. M. Kini, Ed.) *Toxin Reviews*, 40(2), 225-237. doi:10.1080/15569543.2019.1584822
- Hamad, M. T., & Saied, M. S. (2021, January). Kinetic studies of Congo red dye adsorption by immobilized *Aspergillus niger* on alginate. (E. Drioli, Ed.) *Applied Water Science*, 11(2). doi:10.1007/s13201-021-01362-z
- Han, R., Ding, D., Xu, Y., Zou, W., Wang, Y., Li, Y., & Zou, L. (2008, May). Use of rice husk for the adsorption of congo red from aqueous solution in column mode. (A. Pandey, Ed.) *Bioresource Technology*, 99(8), 2938-2946. doi:https://doi.org/10.1016/j.biortech.2007.06.027
- Hassaan, M. A., & Nemr, A. E. (2017, August). Health and Environmental Impacts of Dyes: Mini Review. (M. Minaei, Ed.) *American Journal of Environmental Science and Engineering*, 1(3), 64-67. doi:10.11648/j.ajese.20170103.11
- Hassan, W., Farooq, U., Ahmad, M., Athar, M., & Khan, M. A. (2017, May). Potential biosorbent, *Haloxylon recurvum* plant stems, for the removal of methylene blue dye. (A. A. Alwarthan, Ed.) *Arabian Journal of Chemistry*, 10(2), S1512-S1522. doi:10.1016/j.arabjc.2013.05.002

- Hor, K. Y., Chee, J. M., Chong, M. N., Jin, B., Saint, C., Poh, P. E., & Aryal, R. (2016, April). Evaluation of physicochemical methods in enhancing the adsorption performance of natural zeolite as low-cost adsorbent of methylene blue dye from wastewater. (J. J. Klemeš, C. M. Bôas de Almeida, & Y. Wang, Eds.) *Journal of Cleaner Production*, *118*, 197-209. doi:10.1016/j.jclepro.2016.01.056
- Huang, Y.-H., Tsai, S.-T., Huang, Y.-F., & Chen, C.-Y. (2007, February). Degradation of commercial azo dye reactive Black B in photo/ferrioxalate system. (D. Aga, A. J. Daugulis, Z. He, & G. Lyberatos, Eds.) *Journal of Hazardous Materials*, *140*(1-2), 382-388. doi:10.1016/j.jhazmat.2006.10.083
- Hulme, A. N., Ferreira, E. S., McNab, H., & Quye, A. (2004, February). The natural constituents of historical textile dyes. (J. Love, Ed.) *Chemical Society Reviews*(6), 329-336. doi:10.1039/B305697J
- Ip, A. W., Barford, J. P., & McKay, G. (2010, March). A comparative study on the kinetics and mechanisms of removal of Reactive Black 5 by adsorption onto activated carbons and bone char. (S. Allen, T. (. Aminabhavi, & D. Dionysiou, Eds.) *Chemical Engineering Journal*, *157*(2-3), 434-442. doi:10.1016/j.cej.2009.12.003
- IR Spectrum Table & Chart*. (2022, January 18). Retrieved April 18, 2022, from [www.sigmaaldrich.com](https://www.sigmaaldrich.com): <https://www.sigmaaldrich.com/LK/en/technical-documents/technical-article/analytical-chemistry/photometry-and-reflectometry/ir-spectrum-table>
- Iscen, C. F., Kiran, I., & Ilhan, S. (2007, May). Biosorption of Reactive Black 5 dye by *Penicillium restrictum*: The kinetic study. (D. Aga, A. J. Daugulis, & Z. He, Eds.) *Journal of Hazardous Materials*, *143*(1-2), 335-340. doi:10.1016/j.jhazmat.2006.09.028
- Jeyavishnu, K., & Alagesan, V. (2020, April). *Cereus* sp. as potential biosorbent for removal of Congo red from aqueous solution: isotherm and kinetic

- investigations. (G. B. Wiersma, Ed.) *Environmental Monitoring and Assessment*, 192(4). doi:10.1007/s10661-020-8197-2
- Jia, P., Tan, H., Liu, K., & Gao, W. (2018, August). Removal of Methylene Blue from Aqueous Solution. *Applied science*, 1-11. doi:10.3390/app8101903
- Katheresan, V., Kansedo, J., & Lau, S. Y. (2018, June). Efficiency of Various Recent Wastewater Dye Removal Methods: A Review. *Journal of Environmental Chemical Engineering*, 6(4), 4676-4697. doi:10.1016/j.jece.2018.06.060
- Khadhraoui, M., Trabelsi, H., Ksibi, M., Bouguerra, S., & Elleuch, B. (2009, January). Discoloration and detoxification of a Congo red dye solution by means of ozone treatment for a possible water reuse. (D. Aga, A. J. Daugulis, Z. He, & G. Lyberatos, Eds.) *Journal of Hazardous Materials*, 161(2-3), 974-981. doi:10.1016/j.jhazmat.2008.04.060
- Khan, M. R., Rahman, W., Ong, H. R., Ismail, A. B., & Cheng, C. K. (2016). Tea dust as a potential low-cost adsorbent for the removal of crystal violet from aqueous solution. *Desalination and Water Treatment*, 57(31), 14728-14738. doi:10.1080/19443994.2015.1066272
- Khan, S., Mohammad, A., & Abdul, M. (2019). Mutagenicity and genotoxicity evaluation of textile industry wastewater using bacterial and plant bioassays. (A. M. Tsatsakis, & A. Anadón, Eds.) *Toxicology Reports*, 6, 193-201. doi:10.1016/j.toxrep.2019.02.002
- Khattab, T. A., Abdelrahman, M. S., & Rehan, M. (2020). Textile dyeing industry: environmental impacts and remediation. (P. Garrigues, Ed.) *Environmental Science and Pollution Research*, 27, 3803–3818. doi:10.1007/s11356-019-07137-z
- Krika, F., Krika, A., & Azizi, A. (2019). Arundo donax L. as a low-cost and promising biosorbent for the removal of crystal violet from aqueous media: kinetic, isotherm and thermodynamic investigations. (E. Vessally, Ed.) *Chemical Review and Letters*, 2, 59-68. doi:10.22034/CRL.2019.187434.1015

- Krishna, L. S., Reddy, A. S., Muralikrishna, A., Zuhairi, W. Y., Osman, H., & Reddy, A. V. (2014, September). Utilization of the agricultural waste (Cicer arietinum Linn fruit shell biomass) as biosorbent for decolorization of Congo red. *Desalination and Water Treatment*, 56(8), 2181-2192. doi:10.1080/19443994.2014.958540
- Kristianto, H., Rahman, H., Prasetyo, S., & Sugih, A. K. (2019, May). Removal of Congo red aqueous solution using *Leucaena leucocephala* seed's extract as natural coagulant. (E. Drioli, & S. A. Aljlil, Eds.) *Applied Water Science*, 9(4). doi:10.1007/s13201-019-0972-2
- Kulkarni, M. R., Revanth, T., Acharya, A., & Bhat, P. (2017, March). Removal of Crystal Violet dye from aqueous solution using water hyacinth: Equilibrium, kinetics and thermodynamics study. *Resource-Efficient Technologies*, 3(1), 71-77. doi:10.1016/j.refit.2017.01.009
- Kumar, R., & Ahmad, R. (2011). Biosorption of hazardous crystal violet dye from aqueous solution onto treated ginger waste (TGW). (N. Hilal, Ed.) *Desalination*, 265(1-3), 112-118. doi:10.1016/j.desal.2010.07.040
- Lacasa, E., Cañizares, P., Walsh, F. C., Rodrigo, M. A., & Ponce-de-León, C. (2019, June). Removal of methylene blue from aqueous solutions using an Fe<sup>2+</sup> catalyst and in-situ H<sub>2</sub>O<sub>2</sub> generated at gas diffusion cathodes. (A. R. Hillman, Ed.) *Electrochimica Acta*, 308, 45-53. doi:10.1016/j.electacta.2019.03.218
- Laohaprapanon, S., Matahum, J., Tayo, L., & You, S.-J. (2015, April). Photodegradation of Reactive Black 5 in a ZnO/UV slurry membrane reactor. (H.-K. Tsao, Ed.) *Journal of the Taiwan Institute of Chemical Engineers*, 49, 136-141. doi:10.1016/j.jtice.2014.11.017
- Lellis, B., Polonio, C. Z., Pamphile, J. A., & Polonio, J. C. (2019, December). Effects of textile dyes on health and the environment and bioremediation potential of living organisms. *Biotechnology Research and Innovation*, 3(2), 275-290. doi:10.1016/j.biori.2019.09.001

- Liang, H.-q., Li, M., Liu, B.-t., Xu, Z.-k., & Wu, Q.-y. (2016). Hierarchically Porous Carbon Membranes Derived from PAN and Their Selective Adsorption of Organic Dyes. *Chinese Journal of Polymer Science*, 34(1), 23-33. doi:10.1007/s10118-016-1723-6
- Liang, J., Wu, J., Li, P., Wang, X., & Yang, B. (2012, Feb). Shaddock peel as a novel low-cost adsorbent for removal of methylene blue from dye wastewater. *Desalination and Water Treatment*, 39(1-3), 70-75. doi:10.1080/19443994.2012.669160
- Liao, C.-S., Hung, C.-H., & Chao, S.-L. (2013, February). Decolorization of azo dye reactive black B by *Bacillus cereus* strain HJ-1. (J. d. Boer, T. Galloway, & Y. Yoon, Eds.) *Chemosphere*, 90(7), 2109-2114. doi:10.1016/j.chemosphere.2012.10.077
- Lim, L. B., Priyantha, N., & Mansor, N. H. (2015). Artocarpus altilis (breadfruit) skin as a potential low-cost biosorbent for the removal of crystal violet dye: equilibrium, thermodynamics and kinetics studies. (J. W. LAMOREAUX, Ed.) *Environmental Earth Sciences*, 73(7), 3239–3247. doi:10.1007/s12665-014-3616-8
- Lim, L. B., Usman, A., Hassan, M. H., & Mohamad Zaidi, N. A. (2020). Tropical wild fern (*Diplazium esculentum*) as a new and effective low-cost adsorbent for removal of toxic crystal violet dye. (S. Y. Alraqa, Ed.) *Journal of Taibah University for Science*, 14(1), 621-627. doi:10.1080/16583655.2020.1761122
- Lin, S., Song, Z., Che, G., Ren, A., Li, P., Liu, C., & Zhang, J. (2014, July). Adsorption behavior of metal–organic frameworks for methylene blue from aqueous solution. (W. Schmidt, P. Dietzel, & F. Kleitz, Eds.) *Microporous and Mesoporous Materials*, 139, 27-34. doi:10.1016/j.micromeso.2014.03.004
- Liu, J., Wang, N., Zhang, H., & Baeyens, J. (2019, May). Adsorption of Congo red dye on Fe<sub>x</sub>Co<sub>3-x</sub>O<sub>4</sub> nanoparticles. (R. Dewil, J. Evans, & L. X. Zhang, Eds.)

*Journal of Environmental Management*, 238, 473-483.  
doi:10.1016/j.jenvman.2019.03.009

Lucas, M. S., & Peres, J. A. (2005, September). Decolorization of the azo dye Reactive Black 5 by Fenton and photo-Fenton oxidation. (B. M. Heron, & M. Wainwright, Eds.) *Dyes and Pigments*, 71(3), 236-244.  
doi:10.1016/j.dyepig.2005.07.007

Mahalakshmi, S., Lakshmi, D., & Menaga, U. (2015, Spring). Biodegradation of Different Concentration of dye (Congo red dye) by using Green and Blue Green Algae. (M. Ardestani, & M. H. Niksokhan, Eds.) *International Journal of Environmental Research*, 9(2), 735-744. doi:10.22059/IJER.2015.947

Mahboobeh, D., Hassan, M. E., & Taghi, M. G. (2017). Adsorption Kinetics and Equilibrium Studies of Reactive Red 198 Dye by Cuttlefish Bone Powder. (J. T. Darian, Ed.) *Iranian Journal of Chemistry and Chemical Engineering*, 36(2), 143-151. doi:10.30492/IJCCE.2017.26703

Mahmoodi, N. M., Arami, M., Bahrami, H., & Khorramfar, S. (2011, June). The effect of pH on the removal of anionic dyes from colored textile wastewater using a biosorbent. (S. Spiegel, Ed.) *Applied Polymer Science*, 120(5), 2996–3003. doi:10.1002/app.33406

Malakootian, M., Mansoorian, H. J., Hosseini, A., & Khanjani, N. (2015, July). Evaluating the efficacy of alumina/carbon nanotube hybrid adsorbents in removing Azo Reactive Red 198 and Blue 19 dyes from aqueous solutions. (G. Chen, F. Khan, & M. Hailwood, Eds.) *Process Safety and Environmental Protection*, 96, 125–137. doi:10.1016/j.psep.2015.05.002

Mani, S., & Bharagava, R. N. (2016). Exposure to Crystal Violet, Its Toxic, Genotoxic and Carcinogenic Effects on Environment and Its Degradation and Detoxification for Environmental Safety. *Reviews of Environmental Contamination and Toxicology*, 71-104. doi:10.1007/978-3-319-23573-8\_4

- Mansooreh, D., Medhi, T. M., Talat, G., Mashid, G., Laila, K., Elhameyan, Z., Ghanbarian, M. (2015, April). Optimization of the parameters influencing the photo-fenton process for the decolorization of reactive red 198 (rr198). (M. Ahamadizadeh, Ed.) *Jundishapur journal of health sciences*, 7(2), 38-43. doi:10.5812/jjhs.7(2)2015.28243
- Meriç, S., Kaptan, D., & Ölmez, T. (2004, January). Color and COD removal from wastewater containing Reactive Black 5 using Fenton's oxidation process. (J. d. Boer, Ed.) *Chemosphere*, 54(3), 435-441. doi:10.1016/j.chemosphere.2003.08.010
- Merlot, C., & Bruschiweiler, B. J. (2017, August). Azo dyes in clothing textiles can be cleaved into a series of mutagenic aromatic amines which are not regulated yet. (L. Aylward, & M. v. Berg, Eds.) *Regulatory Toxicology and Pharmacology*, 88, 214-226. doi:10.1016/j.yrtph.2017.06.012
- Mittal, A., Mittal, J., Malviya, A., & Gupta, V. K. (2009, December). Adsorptive removal of hazardous anionic dye "Congo red" from wastewater. (M. Malmsten, Ed.) *Journal of Colloid and Interface Science*, 340(1), 16-26. doi:10.1016/j.jcis.2009.08.019
- Mohamed, R. R., Abu Elell, M. H., Sabaa, M. W., & Saad, G. R. (2018). Synthesis of an efficient adsorbent hydrogel based on biodegradable polymers for removing crystal violet dye from aqueous solution. (A. D. French, Ed.) *Cellulose*, 25(11), 6513–6529. doi:10.1007/s10570-018-2014-x
- Moussavi, G., & Mahmoudi, M. (2009, October). Degradation and biodegradability improvement of the reactive red 198 azo dye using catalytic ozonation with MgO nanocrystals. (S. Allen, T. (. Aminabhavi, & D. Dionysiou, Eds.) *Chemical Engineering Journal*, 152(1), 1-7. doi:10.1016/j.cej.2009.03.014
- Muhammad, U. L., Zango, Z. U., Kadir, H. A., & Usman, A. U. (2019). Crystal Violet removal from aqueous solution using corn stalk biosorbent. (S. G. Abdu, Ed.) *Science World Journal*, 14(1), 133-138. Retrieved from <https://www.ajol.info/index.php/swj/article/view/208516>

- Munagapati, S. V., & Kim, D. S. (2016, August). Adsorption of anionic azo dye Congo Red from aqueous solution. (W. Schröer, & T. Yamaguchi, Eds.) *Journal of Molecular Liquids*, 220, 540-548. doi:10.1016/j.molliq.2016.04.119
- Munagapati, V. S., Wen, J.-C., Pan, C.-L., Gutha, Y., Wen, J.-H., & Reddy, G. M. (2019, August). Adsorptive removal of anionic dye (Reactive Black 5) from aqueous solution using chemically modified banana peel powder: kinetic, isotherm, thermodynamic, and reusability studies. (L. A. Newman, Ed.) *International Journal of Phytoremediation*, 22(3), 267-278. doi:10.1080/15226514.2019.1658709
- Munagapati, V. S., Yarramuthi, V., Kim, Y., Lee, K. M., & Kim, D.-S. (2018, February). Removal of anionic dyes (Reactive Black 5 and Congo Red) from aqueous solutions using Banana Peel Powder as an adsorbent. (R. Handy, & B. Yan, Eds.) *Ecotoxicology and Environmental Safety*, 148, 601-607. doi:10.1016/j.ecoenv.2017.10.075
- Nandiyanto, A. B., Oktiani, R., & Ragadhita, R. (2019). How to Read and Interpret FTIR Spectroscopy of Organic Material. *Indonesian Journal of Science & Technology*, 97-118. doi:10.17509/ijost.v4i1.15806
- Neupane, S., Ramesh, S. T., Gandhimathi, R., & Nidheesh, P. V. (2015). Pineapple leaf (*Ananas comosus*) powder as a biosorbent for the removal of crystal violet from aqueous solution. *Desalination and Water Treatment*, 54(7), 2041-2054. doi:10.1080/19443994.2014.903867
- Nikfar, S., & Jaberidoost, M. (2014). Dyes and Colorants. In P. Wexler (Ed.), *Encyclopedia of Toxicology* (3 ed., Vol. 2, pp. 252-261). Academic Press. doi:10.1016/B978-0-12-386454-3.00602-3
- Ooi, J. Q., Morad, N., & Tan, A. K. (2016, October). Phytoremediation of Methylene Blue and Methyl Orange Using *Eichhornia crassipes*. (R. Haynes, Ed.) *International Journal of Environmental Science and Development*, 7(10), 724-728. doi:10.18178/ijesd.2016.7.10.869



- Orozco, R. S., Martínez-Juan, M., García-Sánchez, J. J., & Ureña-Núñez, F. (2018, January). Removal of Methylene Blue from Aqueous Solution Using Typha Stems and Leaves. (L. Lucia, & M. Hubbe, Eds.) *BioResources*, *13*(1), 1696-1710. doi:10.15376/biores.13.1.1696-1710
- Osma, J. F., Saravia, V., Toca-Herrera, J. L., & Couto, S. R. (2007, August). Sunflower seed shells: A novel and effective low-cost adsorbent for the removal of the diazo dye Reactive Black 5 from aqueous solutions. (D. Aga, A. J. Daugulis, & Z. He, Eds.) *Journal of Hazardous Materials*, *147*(3), 900-905. doi:10.1016/j.jhazmat.2007.01.112
- Pal, P. (2017). Industry-Specific Water Treatment: Case Studies. In P. Pal, *Industrial Water Treatment Process Technology* (1 ed., pp. 243-511). Butterworth-Heinemann. doi:10.1016/B978-0-12-810391-3.00006-0
- Palma-Goyes, R. E., Guzmán-Duque, F. L., Peñuela, G., González, I., Nava, J. L., & Torres-Palma, R. A. (2010, September). Electrochemical degradation of crystal violet with BDD electrodes: Effect of electrochemical parameters and identification of organic by-products. (J. de Boer, T. Galloway, & Y. Yoon, Eds.) *Chemosphere*, *81*(1), 26-32. doi:10.1016/j.chemosphere.2010.07.020
- Patel, H., & Vashi, R. T. (2012, April). Removal of Congo Red dye from its aqueous solution. (A. A. Mayouf, Ed.) *Journal of Saudi Chemical Society*, *16*(2), 131-136. doi:10.1016/j.jscs.2010.12.003
- Pathania, D., Sharma, A., & Siddiqi, Z.-M. (2016, July). Removal of congo red dye from aqueous system using Phoenix dactylifera seeds. (W. Schröer, T. Yamaguchi, & A. J. Valente, Eds.) *Journal of Molecular Liquids*, *219*, 359-367. doi:10.1016/j.molliq.2016.03.020
- Paz, D. S., Baiotto, A., Schwaab, M., Mazutti, M. A., Bassaco, M. M., Bertuol, D. A., Meili, L. (2013, July). Use of papaya seeds as a biosorbent of methylene blue from aqueous solution. (W. Rauch, Ed.) *Water Science & Technology*, *68*(2), 441–447. doi:10.2166/wst.2013.185

- Pimentel, P., Oliveira, R., Melo, D., Melo, M., Assunção, A., & Gonzales, G. (2011). Adsorption of chromium ions on oil shale waste. *Brazilian journal of petroleum and gas*, 5(2), 065-073. doi:10.5419/bjpg2011-0008
- Qingxiang, Y., Lingxia, T., Min, Y., & Hao, Z. (2007, June). Effects of glucose on the decolorization of Reactive Black 5 by yeast isolates. (J. Hao, X. Le, & W. Mitch, Eds.) *Journal of Environmental Sciences*, 20(1), 105-108. doi:10.1016/S1001-0742(08)60016-9
- Qureshi, U. A., Khatri, Z., Ahmed, F., Khatri, M., & Kim, I.-S. (2017, April). Electrospun Zein Nanofiber as a Green and Recyclable Adsorbent for the Removal of Reactive Black 5 from the Aqueous Phase. (D. T. Allen, Ed.) *ACS Sustainable Chemistry & Engineering*, 5(5), 4340–4351. doi:10.1021/acssuschemeng.7b00402
- Rahimi, s., Ehrampoush, m. h., Ghaneian, m. t., Reshadat, s., fatehizadeh, a., Ahmadian, m., Rahimi, s. (2013, October). Application of TiO<sub>2</sub>/UV-C Photocatalytic Process in Removal of Reactive Red 198 Dye from Synthetic Textile Wastewater. *Asian Journal of Chemistry*, 25(13), 7427-7432. doi:10.14233/ajchem.2013.14764
- Rani, K. C., Naik, A., Chaurasiya, R. S., & Raghavarao, K. (2017, May). Removal of toxic Congo red dye from water employing low-cost coconut residual fiber. (W. Rauch, Ed.) *Water Science & Technology*, 75(9). doi:10.2166/wst.2017.109
- Raymundo, A. S., Zanarotto, R., Belisário, M., Pereira, M. d., Ribeiro, J. N., & Ribeiro, A. V. (2010, August). Evaluation of Sugar-Cane Bagasse as Bioadsorbent in the Textile Wastewater Treatment Contaminated with Carcinogenic Congo Red Dye. *Brazilian Archives of Biology and Technology*, 53(4), 931-938. doi:10.1590/S1516-89132010000400023
- REACTIVE DYES.** (2021, August 20). (Global Colors, Inc) Retrieved July 10, 2021, from Global Colors Inc: <https://globalcolorsinc.com/reactive-dyes/>

- Reactive red 198*. (n.d.). Retrieved 9 2021, from ChemicalBook.com:  
[https://www.chemicalbook.com/ChemicalProductProperty\\_EN\\_CB9260610.htm](https://www.chemicalbook.com/ChemicalProductProperty_EN_CB9260610.htm)
- reactive-black-5*. (2012, May 18). Retrieved from worlddyevariety:  
<http://www.worlddyevariety.com/reactive-dyes/reactive-black-5.html>
- Ruthiraan, M., Abdullah, E. C., Mubarak, N. M., & Noraini, M. N. (2017, April). A promising route of magnetic based materials for removal of cadmium and methylene blue from waste water. (G. L. Dotto, & D. Fatta-Kassinos, Eds.) *Journal of Environmental Chemical Engineering*, 5(2), 1447-1455. doi:10.1016/j.jece.2017.02.038
- S, R., Lata, S., & P, B. (2018, March). Biosorption characteristics of methylene blue and malachite green from simulated wastewater onto Carica papaya wood biosorbent. (M. Chipara, H. Deng, & J. Gao, Eds.) *Surfaces and Interfaces*, 10, 197-215. doi:10.1016/j.surfin.2017.09.011
- Saha, P. (2010). Assessment on the Removal of Methylene Blue Dye using Tamarind Fruit Shell as Biosorbent. (J. T. Trevors, Ed.) *Water, Air, & Soil Pollution*, 213(1-4), 287–299. doi:10.1007/s11270-010-0384-2
- Sahoo, C., Gupta, A. K., & Pal, A. (2005, September). Photocatalytic degradation of Crystal Violet (C.I. Basic Violet 3) on silver ion doped TiO<sub>2</sub>. (B. M. Heron, & M. Wainwright, Eds.) *Dyes and Pigments*, 66(3), 189-196. doi:10.1016/j.dyepig.2004.09.003
- Saini, R. D. (2017). Textile Organic Dyes: Polluting effects and Elimination Methods from Textile Waste Water. (R. Gupta, Ed.) *International Journal of Chemical Engineering Research*, 9, 121-136. Retrieved from [http://www.ripublication.com/ijcher17/ijcherv9n1\\_10.pdf](http://www.ripublication.com/ijcher17/ijcherv9n1_10.pdf)
- Salehi, M., Hashemipour, H., & Mirzaee, M. (2012). Experimental Study of Influencing Factors and Kinetics in Catalytic Removal of Methylene Blue with TiO<sub>2</sub> Nanopowder. (S. Bonacci, Ed.) *American Journal of Environmental Engineering*, 2(1), 1-7. doi:10.5923/j.ajee.20120201.01

- Salem, E. A., Fayed, T. A., El-Nahass, M. N., & Dawood, M. (2018, January). A Comparative Study for Adsorption of Methylene Blue Dye from Wastewater on to Three Different Types of Rice Ash. (A. El-Saghier, Ed.) *Journal of Pharmaceutical and Applied Chemistry*, 4(2), 99-107. doi:10.18576/jpac/040204
- Sarvestani, M. R., & Doroudi, Z. (2020). Removal of Reactive Black 5 from Waste Waters by Adsorption: A Comprehensive Review. (S. A. Bidgoli, & M. Jahanshahi, Eds.) *Journal of Water and Environmental Nanotechnology*, 5(2), 180-190. doi:10.22090/JWENT.2020.02.008
- Sathishkumar, K., AlSalhi, M. S., Sanganyado, E., Devanesan, S., Arulprakash, A., & Rajasekar, A. (2019, November). Sequential electrochemical oxidation and bio-treatment of the azo dye congo red and textile effluent. (A. L. Pinheiro, D. Ramaiah, & M. d. Baptista, Eds.) *Journal of Photochemistry and Photobiology B: Biology*, 200. doi:10.1016/j.jphotobiol.2019.111655
- Sekar, N. (2011). Acid dyes. In M. Clark (Ed.), *Handbook of Textile and Industrial Dyeing: Principles, Processes and Types of Dyes* (1 ed., Vol. 1, pp. 486-514). Woodhead Publishing. doi:10.1533/9780857093974.2.486
- Selim, K. A., Abdel-Khalek, N. A., El-Sayed, S. M., & Abdallah, S. S. (2015, Nvember). Bioremoval of Crystal Violet Dye from Egyptian Textile Effluent. *International Research Journal of Engineering and Technology (IRJET)*, 2(8), 1038-1043. Retrieved from <https://www.irjet.net/archives/V2/i8/IRJET-V2I8155.pdf>
- Sharma, S., Hasan, A., Kumar, N., & Pandey, L. M. (2018, May). Removal of methylene blue dye from aqueous solution using immobilized *Agrobacterium fabrum* biomass along with iron oxide nanoparticles as biosorbent. (P. Garrigues, Ed.) *Environmental Science and Pollution Research*, 25(22), 21605–21615. doi:10.1007/s11356-018-2280-z
- Shirzad-Siboni, M., Jafari, S. J., Giahi, O., Kim, I., Lee, S.-M., & Yang, J.-K. (2014, July). Removal of acid blue 113 and reactive black 5 dye from aqueous

- solutions by activated red mud. (B. R. Yoo, Ed.) *Journal of Industrial and Engineering Chemistry*, 20(4), 1432-1437. doi:10.1016/j.jiec.2013.07.028
- Silva, L. S., Lima, L. C., Ferreira, F. J., Silva, M. S., Osajima, J. A., Bezerra, R. D., & Filho, S. E. (2015, April). Sorption of the anionic reactive red RB dye in cellulose: Assessment of kinetic, thermodynamic, and equilibrium data. (J. Plumet, Ed.) *Open Chemistry*, 13(1), 801-812. doi:10.1515/chem-2015-0079
- Singhapon , T., Shinoda , K., Rujakom, S., & Kazama, F. (2021, March). Factors Affecting the Simultaneous Removal of Nitrate and Reactive Black 5 Dye via Hydrogen-Based Denitrification. (J.-L. PROBST, Ed.) *Water*, 13(7). doi:10.3390/w13070922
- Stjepanović, M., Velić, N., Galić , A., Kosović , I., Jakovljević , T., & Stanić, M. H. (2021, February). From Waste to Biosorbent: Removal of Congo Red from Water by Waste Wood Biomass. (M. M. Manca, Ed.) *Water*, 13(3). doi:10.3390/w13030279
- Surana, M., Mehta, P., Pamecha, K., & Kabra, B. V. (2011). Treatment of water contaminated with Reactive Red 198 (RR198) by Photo-Fenton Reagent. *Pelagia Research Library*, 2(2), 177-186. Retrieved from <https://www.imedpub.com/articles/treatment-of-water-contaminated-with-reactive-red-198-rr198by-photofenton-reagent.pdf>
- Tan , I. A., & Hameed, B. H. (2012). Removal of crystal violet dye from aqueous solutions using rubber (hevea brasillensis) seed shell-based biosorbent. *Desalination and Water Treatment*, 48(1-3), 174-181. doi:10.1080/19443994.2012.698810
- Tang, C., & Chen, V. (2004, June). The photocatalytic degradation of reactive black 5 using TiO<sub>2</sub>/UV in an annular photoreactor. (E. Morgenroth, G.-H. Chen, & A. Deletic, Eds.) *Water Research*, 38(11), 2775-2781. doi:10.1016/j.watres.2004.03.020
- Tayebi, H.-A., Dalirandeh, Z., Rad, A. S., Mirabi, A., & Binaeian, E. (2016, January). Synthesis of polyaniline/Fe<sub>3</sub>O<sub>4</sub> magnetic nanoparticles for removal

of reactive red 198 from textile waste water: kinetic, isotherm, and thermodynamic studies. *Desalination and Water Treatment*, 57(47), 22551-22563. doi:10.1080/19443994.2015.1133323

*Technical report 123 Freundlich isotherms*. (n.d.). Retrieved from ecetoc: <https://www.ecetoc.org/report/measured-partitioning-property-data/adsorption-desorption-distribution-kd-and-organic-carbon-water-partition-koc-coefficients/freundlich-isotherms/>

Tichapondwa, S., Newman, J., & Kubheka, O. (2020, October). Effect of TiO<sub>2</sub> phase on the photocatalytic degradation of methylene blue dye. (J. Gottsmann, J. Liu, & M. Mul, Eds.) *Physics and Chemistry of the Earth, Parts A/B/C*, 118-119. doi:10.1016/j.pce.2020.102900

Ullah, I., Ali, S., & Akram, M. (2013). Degradation of Reactive Black B dye in wastewater using oxidation process. (M. A. Hanif, Ed.) *International Journal of Chemical and Biochemical Sciences*, 4, 96-100. doi:10.1.1.707.5470

UrRehman, M. S., Kim, I., & Han, J.-I. (2012, October). Adsorption of methylene blue dye from aqueous solution by sugar extracted spent rice biomass. (M. Coimbra, K. Edgar, & T. Budtova, Eds.) *Carbohydrate Polymers*, 90(3), 1314-1322. doi:10.1016/j.carbpol.2012.06.078

Vargas, V. H., Paveglio, R. R., Pauletto, P., Gonçalves Salau, N., & Dotto, L. (2019, April). Sisal fiber as an alternative and cost-effective adsorbent for the removal of methylene blue and reactive black 5 dyes from aqueous solutions. (A. Gill, Ed.) *Chemical Engineering Communications*, 207(4), 523-536. doi:10.1080/00986445.2019.1605362

Vijayaraghavan, G., & Shanthakumar, S. (2015, February). Performance study on algal alginate as natural coagulant for the removal of Congo red dye. *Desalination and Water Treatment*, 57(14), 6384-6392. doi:10.1080/19443994.2015.1008578

Wanyonyi, W. C., Onyari, J. M., & Shiundu, P. M. (2014, July). Adsorption of Congo Red Dye from Aqueous Solutions Using Roots of Eichhornia

- Crassipes: Kinetic and Equilibrium Studies. *Energy Procedia*, 50, 862-869. doi:10.1016/j.egypro.2014.06.105
- Wathukarage, A., Herath, I., Iqbal, M. C., & Vithanage, M. (2019). Mechanistic understanding of crystal violet dye sorption by woody biochar: implications for wastewater treatment. (M. H. Wong, Ed.) *Environmental Geochemistry and Health*, 1647–1661. doi:10.1007/s10653-017-0013-8
- Wathukarage, A., Herath, I., Iqbal, M., & Vithanage, M. (2017, August). Mechanistic understanding of crystal violet dye sorption by woody biochar: implications for wastewater treatment. *Environ Geochem Health*. doi:10.1007/s10653-017-0013-8
- Wekoye, J. N., Wanyonyi, W. C., Wangila, P. T., & Tonui, M. K. (2020). Kinetic and equilibrium studies of Congo red dye adsorption on cabbage waste powder. (K. Kannan, Ed.) *Environmental Chemistry and Ecotoxicology*, 2, 24-31. doi:10.1016/j.eneco.2020.01.004
- Won, S. W., Kim, H.-J., Choi, S.-H., Chung, B.-W., Kim, K.-J., & Yun, Y.-S. (2006, August). Performance, kinetics and equilibrium in biosorption of anionic dye Reactive Black 5 by the waste biomass of *Corynebacterium glutamicum* as a low-cost biosorbent. (S. Allen, T. Aminabhavi, & D. Dionysiou, Eds.) *Chemical Engineering Journal*, 101(1), 37-43. doi:10.1016/j.cej.2006.04.005
- Xia, H., Chen, L., & Fang, Y. (2013, August). Highly Efficient Removal of Congo red from Wastewater by Nano-Cao. (R. Wickramasinghe, Ed.) *Separation Science and Technology*, 48(17), 2681-2687. doi:10.1080/01496395.2013.805340
- Yang, Y., Ali, N., Khan, A., Khan, S., Khan, S., Khan, H., Bilal, M. (2021, January). Chitosan-capped ternary metal selenide nanocatalysts for efficient degradation of Congo red dye in sunlight irradiation. (A. Dong, & J. F. Kennedy, Eds.) *International Journal of Biological Macromolecules*, 167, 169-181. doi:10.1016/j.ijbiomac.2020.11.167

- You, S.-J., Damodar, R. A., & Hou, S.-C. (2010, May). Degradation of Reactive Black 5 dye using anaerobic/aerobic membrane bioreactor (MBR) and photochemical membrane reactor. (D. Aga, A. J. Daugulis, & Z. He, Eds.) *Journal of Hazardous Materials*, 173(1-3), 1112-1118. doi:10.1016/j.jhazmat.2010.01.036
- Zahir, A., Aslam, Z., Kamal, M. S., Ahmad, W., Abbas, A., & Shawabkeh, R. A. (2017, October). Development of novel cross-linked chitosan for the removal of anionic Congo red dye. (W. Schroer, T. Yamaguchi, & A. J. Valente, Eds.) *Journal of Molecular Liquids*, 244, 211-218. doi:10.1016/j.molliq.2017.09.006
- Zamora, M. H., Jeronimo, F. M., Urbina, E. C., & Villanueva, R. O. (2016, September). Congo red dye affects survival and reproduction in the cladoceran *Ceriodaphnia dubia*. Effects of direct and dietary exposure. (L. R. Shugart, Ed.) *Ecotoxicology*, 25(10), 1832-1840. doi:10.1007/s10646-016-1731-x
- Zare, K., Sadegh, H., ghoshekandi, R. S., Maazinejad, B., Ali, V., Tyagi, I., Gupta, V. K. (2015, December). Enhanced removal of toxic Congo red dye using multi walled carbon nanotubes: Kinetic, equilibrium studies and its comparison with other adsorbents. (W. Schröer, T. Yamaguchi, A. J. Valente, & L. F. Vega, Eds.) *Journal of Molecular Liquids*, 212, 266-271. doi:10.1016/j.molliq.2015.09.027
- Zheng, H., Sun, Q., Li, Y., & Du, Q. (2020, April). Biosorbents prepared from pomelo peel by hydrothermal technique and its adsorption properties for congo red. (A. Balazs, & S. Banerjee, Eds.) *Materials Research Express*, 7. doi:10.1088/2053-1591/ab8a83